

A linear-time algorithm to compute total $[1, 2]$ -domination number of block graphsPouyeh Sharifani^{a,c}, Mohammadreza Hooshmandasl^{*b,c}, Saeid Alikhani^d^aInstitute for Research in Fundamental Sciences (IPM), Tehran, Iran^bDepartment of Computer Science, University of Mohaghegh Ardabili, Ardabil, Iran^cDepartment of Computer Science, Yazd University, Yazd, Iran^dDepartment of Mathematics, Yazd University, Yazd, Iran

ABSTRACT: Let $G = (V, E)$ be a simple graph without isolated vertices. A set $D \subseteq V$ is a total $[1, 2]$ -dominating set if for every vertex $v \in V$, $1 \leq |N(v) \cap D| \leq 2$. The total $[1, 2]$ -domination problem is to determine the total $[1, 2]$ -domination number $\gamma_{t[1,2]}(G)$, which is the minimum cardinality of a total $[1, 2]$ -dominating set for a graph G . In this paper, we present a linear-time algorithm to compute $\gamma_{t[1,2]}(G)$ for a block graph G .

Review History:

Received:19 May 2020

Revised:10 August 2020

Accepted:14 August 2020

Available Online:01 September 2020

Keywords:Total $[1, 2]$ -set
Dominating set
Block graph**AMS Subject classifications:**

05C15; 20D60

1. Introduction

All graphs considered here are simple, i.e., finite, undirected, and loop-less. For other graph theory terminology and notation not given here we refer to [10].

Let $G = (V, E)$ be a graph. The *open neighborhood* of a vertex $v \in V$ is the set of all vertices adjacent to v and is denoted by $N(v)$. Similarly, the *closed neighborhood* of a vertex v is $N[v] = N(v) \cup \{v\}$. In connected graph G , a vertex is called a *cut-vertex* of G if its removal produces a disconnected graph. A *block of a graph* G is a maximal connected induced subgraph of G that has no cut-vertex. A block graph is a graph whose blocks are complete graphs. A subset $D \subseteq V$ is called a *dominating set*, if every vertex in V is contained in D or has a neighbor in D . The domination number of G , denoted by $\gamma(G)$, is the minimum cardinality of a dominating set. A *total dominating set* of a simple graph $G = (V, E)$ without isolated vertex, is a set $S \subseteq V$ such that every vertex is adjacent to a vertex in S . The minimum cardinality of a total dominating set is denoted by $\gamma_t(G)$. The minimum dominating set problem is an *NP-hard* problem [8, 7, 9].

A set $S \subseteq V$ is called a $[1, 2]$ -set of G if for each $v \in V - S$, v is adjacent to at least one but not more than two vertices in S . The total $[1, 2]$ -set is a set $S \subseteq V$ such that for each $v \in V$, $1 \leq |N(v) \cap S| \leq 2$. The total $[1, 2]$ -domination number, denoted by $\gamma_{t[1,2]}(G)$, is the minimum cardinality of a total $[1, 2]$ -dominating set for a graph G . We note that in the problem total $[1, 2]$ -dominating set, there are some graphs without any total $[1, 2]$ -dominating sets, such as the graphs with isolated vertices.

*Corresponding author.

E-mail addresses: pouyeh.sharifani@gmail.com, hooshmandasl@uma.ac.ir, hooshmandasl@yazd.ac.ir, alikhani@yazd.ac.ir

The concept of $[1, 2]$ -set and its variants such as total $[1, 2]$ -set and independent $[1, 2]$ -set are well studied by Chellali et al. in [3, 4]. In [4], several open problems were proposed about $[1, 2]$ -set and total $[1, 2]$ -set. Some of these problems were solved in [6, 2, 5]. In this paper we provide a linear-time algorithm for finding the minimum cardinality of a total $[1, 2]$ -dominating set for a block graph G .

2. Algorithm for computing total $[1, 2]$ -domination number

Our algorithm relies on a tree-like decomposition structure, which is called a refined cut-tree of a block graph.

Let G be a block graph with t blocks B_1, B_2, \dots, B_t and q cut-vertices v_1, v_2, \dots, v_q . The cut-tree of G , denoted by $T^C(V^C, E^C)$, is defined as $V^C = \{B_1, B_2, \dots, B_t, v_1, v_2, \dots, v_q\}$ and $E^C = \{(B_i, v_j) \mid v_j \in B_i, 1 \leq i \leq h, 1 \leq j \leq q\}$. The cut-tree of a block graph can be constructed in linear-time by the depth-first search algorithm [1]. For any block B_i of G , the block-vertex \tilde{B}_i is defined as $\tilde{B}_i = \{v \in B_i \mid v \text{ is not a cut-vertex}\}$, where $1 \leq i \leq t$. We can refine the cut-tree $T^C(V^C, E^C)$ as $V^C = \{\tilde{B}_1, \dots, \tilde{B}_t, v_1, \dots, v_q\}$ and $E^C = \{(\tilde{B}_i, v_j) \mid v_j \in B_i, 1 \leq i \leq t, 1 \leq j \leq q\}$. We notice that in the refined cut-tree of a block graph, a block-vertex can be empty.

A block graph G with 11 blocks B_1, B_2, \dots, B_{11} and the corresponding refined cut-tree of G are shown in Figure 1.




Figure 1: Block graph G and the corresponding cut-tree of G

To compute $\gamma_{[1,2]}(G)$, we traverse T in the post order and during traversing, we compute $m_i^-[v]$, $m_i^+[v]$, $S_0^-(u)$, $S_{0,1}^-(u)$, $S_{1,2}^-(u)$, $S_0^+(u)$, $S_{0,1}^+(u)$ and $S^*(u)$ where $i = 0, 1, 2$, v is cut-vertex and $u \in V(T)$ is block node.

- B as the set of all block nodes of T .
- C as the set of all cut-vertices of G .
- T_v as the subtree of T rooted at v .
- $G[T_v]$ as the subgraph of G which corresponding to T_v .
- For the smallest $[1, 2]$ -set S of T_v , every vertex in S is called (S, v) -black, and (S, v) -white otherwise. For simplicity, we use the terms black and white instead of (S, v) -black and (S, v) -white, respectively.
- For each block node $u \in V(T)$ and $v \in ch(u)$, depended on the color of v and the number of vertices which dominate v , we define variables as bellow:
 - For $u \in V(T)$, $S_0^-(u)$ is the size of the smallest $[1, 2]$ -set of $G[T_v]$ that all children of u are white and they are not dominated.
 - For $u \in V(T)$, $S_{0,1}^-(u)$ is the size of the smallest $[1, 2]$ -set of $G[T_v]$ that all children of u are white and they are dominated at most once.
 - For $u \in V(T)$, $S_{1,2}^-(u)$ is the size of the smallest $[1, 2]$ -set of $G[T_v]$ that all children of u are white and they are dominated once or twice.
 - For $u \in V(T)$, $S_0^+(u)$ is the size of the smallest $[1, 2]$ -set of $G[T_v]$ that one child of u is black and it is dominated at most once. Moreover the other children of u are white and they are not dominated.

- For $u \in V(T)$, $S_{0,1}^+(u)$ is the size of the smallest $[1, 2]$ -set of $G[T_v]$ that one vertex of $ch(u)$ is black and it is dominated once or twice. Moreover the other children of u are white and they are dominated at most once.
- For $u \in V(T)$, $S^*(u)$ is the size of the smallest $[1, 2]$ -set of $G[T_v]$ that two black vertices of $ch(u)$ are dominated at most once. Moreover the other children of u are white and they are not dominated.
- For cut-vertex $v \in V(T)$, $m_i^-[v]$ is the size of the smallest $[1, 2]$ -set of $G[T_v]$ that v is white and dominated by i other vertices of $G[T_v]$ for $i = 0, 1, 2$.
- For cut-vertex $v \in V(T)$, $m_i^+[v]$ is the size of the smallest $[1, 2]$ -set of $G[T_v]$ that v is black and dominated by i other vertices of $G[T_v]$ for $i = 0, 1, 2$.

Now, we use a refined cut tree T of a given block graph G and dynamic programming method to compute total $[1, 2]$ -domination number of G . The algorithm contains three step, Initializing step, Updating step and final step.

2.1. Initializing step:

Obviously, every leaf of cut-tree T is a block node and it is not empty. Since variables for block nodes are based on color of its child, so in first step we begin our algorithm from pre-pendent node v of T , that is a cut vertex of G . We initialize $m_i^+[v]$ and $m_i^-[v]$ for $i = 0, 1, 2$ and pre-pendent node v of T as bellow:

$$m_0^+[v] = 1, \quad m_1^+ = 2, \quad m_2^+ = 3, \quad m_0^-[v] = \infty,$$

$$m_1^-[v] = \begin{cases} 1 & \text{if } |ch(v)| = 1, \\ \infty & \text{Otherwise,} \end{cases} \quad \text{and} \quad m_2^-[v] = \begin{cases} 2 & \text{if } |ch(v)| = 2, \\ \infty & \text{Otherwise.} \end{cases}$$

2.2. Updating step:

In the post order traversal of T , for each non pre-pendent node, based on type of them which are a block or cut, we can consider the following cases:

2.3. Updating step for block nodes of T :

In this step, we define variables $S_0^-(u), S_{0,1}^-(u), S_{1,2}^-(u), S_0^+(u), S_{0,1}^+(u)$ and $S^*(u)$ for block node u of T . These variables depend on the number of nodes in $ch(u)$.

Calculating $S_0^-(u)$:

All children of u are white and they are not dominated. So:

$$S_0^-(u) = \sum_{v \in ch(u)} m_0^-(v).$$

Calculating $S_{0,1}^-(u)$:

All children of u are white and they are dominated at most once. So:

$$S_{0,1}^-(u) = \sum_{v \in ch(u)} \text{Min}\{m_0^-(v), m_1^-(v)\}.$$

Calculating $S_{1,2}^-(u)$:

All children of u are white and they are dominated at most once. So:

$$S_{1,2}^-(u) = \sum_{v \in ch(u)} \text{Min}\{m_1^-(v), m_2^-(v)\}.$$

Calculating $S_0^+(u)$:

One child $v_i \in ch(u)$ is black, it is dominated at most once and the other children of u are white and they are not dominated. So:

$$S_0^+(u) = \text{Min}_{v_i \in ch(u)} \{ \text{Min}\{m_0^+(v_i), m_1^+(v_i)\} + \sum_{v \in ch(u), v \neq v_i} m_0^-(v) \}.$$

Calculating $S_{0,1}^+(u)$:

One child $v_i \in ch(u)$ is black and it is dominated once or twice. Moreover the other children of u are white and they are dominated at most once. So:

$$S_{0,1}^+(u) = \text{Min}_{v_i \in ch(u)} \{ \text{Min} \{ m_1^+(v_i), m_2^+(v_i) \} + \sum_{v \in ch(u), v \neq v_i} \text{Min} \{ m_0^-(v), m_1^-(v) \} \}.$$

Calculating $S^*(u)$:

Two vertices of $v_i, v_{i'} \in ch(u)$ are black and they are dominated at most once. Moreover the other children of u are white and they are not dominated. So:

$$S^*(u) = \text{Min}_{v_i, v_{i'} \in ch(u)} \{ \text{Min} \{ m_0^+(v_i), m_1^+(v_i) \} + \{ \text{Min} \{ m_0^+(v_{i'}), m_1^+(v_{i'}) \} + \sum_{v \in ch(u), v \neq v_i, v_{i'}} m_0^-(v_i) \}.$$

2.4. Updating step for cut vertex of T :

In this step, for $i = 0, 1, 2$ we define variables $m_0^-i(v)$ and $m_i^+(v)$ for cut nodes v of T .

Calculating $m_0^+(v)$ when v is not a pre-pendent:

In this case, v is black so all the children of v and all children of $ch(v)$ already have a black neighbor. Since v should not dominate by any vertices, So, all child $u \in ch(v)$ must be white and they are dominated at most once.

$$m_0^+(v) = 1 + \sum_{u \in ch(v)} S_{0,1}^-(u).$$

Calculating $m_1^+(v)$ when v is not a pre-pendent:

In this case, v is black and it is dominated once, so one of the following cases can occurs:

- In this case, exactly one vertex of block $u_i \in ch(v)$ is black and other children of u_i have color white and are not dominated. For the other block $u \in ch(v)$, all child are white and they are dominated at most once. We have:

$$M_1^+ = 1 + \text{Min}_{u_i \in ch(v), |u_i| \neq 0} \{ S_0^-(u_i) + \sum_{u \in ch(v), u \neq u_i} S_{0,1}^-(u) \}.$$

- For exactly one block $u_i \in ch(v)$, one nodes of $ch(u_i)$ and the black vertex dominate at most once. The other vertices of $ch(u_i)$ are white and they are not dominated. In addition, for the other block $u \in ch(v)$, all of their children are white and they are dominated at most once. We have:

$$M_1'^+ = \text{Min}_{u_i \in ch(v)} \{ S_0^+(u_i) + \sum_{u \in ch(v), u \neq u_i} S_{0,1}^-(u) \}.$$

Minimum of M_1^+ and $M_1'^+$ is the best value for $m_1^+(v)$. So:

$$m_1^+(v) = \text{Min} \{ M_1^+, M_1'^+ \}.$$

Calculating $m_2^+(v)$ when v is not a pre-pendent:

In this case, v is black and it is dominated twice, so one of the following cases can occurs:

- There exist at least one vertex $u_i \in ch(v)$ such that $|u_i| = |ch(u_i)| = 1$. In this case, vertex in block u_i is black, its child is black and it is not dominated. So we have:

$$M_2^+ = 1 + \text{Min}_{u_i \in ch(v), |u_i|=|ch(u_i)|=1} \{ m_0^+(ch(u_i)) \} + \sum_{u \in ch(v), u \neq u_i} S_{0,1}^-(u).$$

If there is not any node $u_i \in ch(v)$ such that $|u_i| = |ch(u_i)| = 1$, then $M_2^+ = \infty$.

- There exist at least one vertex $u_i \in ch(v)$ such that $|u_i| = 0$ and $|ch(u_i)| = 2$. In this case, both children of $ch(u_i)$ are black and it is not dominated. So we have:

$$M_2'^+ = \text{Min}_{u_i \in ch(v), |u_i|=0, |ch(u_i)|=2} \{ m_0^+(ch(u_i)) \} + \sum_{u \in ch(v), u \neq u_i} S_{0,1}^-(u).$$

If there is not any node $u_i \in ch(v)$ such that $|u_i| = |ch(u_i)| = 1$, then $M_2'^+ = \infty$.

- For exactly two block $u_i, u_j \in ch(v)$, one vertex is black, so all children of $ch(u_i)$ and $ch(u_j)$ should be white and they are not dominated. For the other block $u \in ch(v)$, all child must be white and they are dominated at most once. We have:

$$M_2''^+ = 2 + \text{Min}_{u_i, u_j \in ch(v)} \{S_0^-(u_i) + S_0^-(u_j) + \sum_{u \in ch(v), u \neq u_i, u_j} S_{0,1}^-(u)\}.$$

- For exactly one block $u_i \in ch(v)$, one vertex is black, so all children of $ch(u_i)$ should be white and they are not dominated. In addition, for exactly one block $u_j \in ch(v)$, exactly one child is black and dominated at most once and the other child is white and not dominated. Also, all children other child $u \in ch(v)$ should be white and they are dominated at most once.. We have:

$$M_2'''^+ = 1 + \text{Min}_{u_i, u_j \in ch(v)} \{S_0^-(u_i) + S_0^+(u_j) + \sum_{u \in ch(v), u \neq u_i, u_j} S_{0,1}^-(u)\}.$$

- For exactly two blocks $u_i, u_j \in ch(v)$, exactly one child is black and dominated at most once and the other child is white and not dominated. Also, all children other child $u \in ch(v)$ should be white and they are dominated at most once.. We have:

$$M_2''''^+ = \text{Min}_{u_i, u_j \in ch(v)} \{S_0^-(u_i) + S_0^-(u_j) + \sum_{u \in ch(v), u \neq u_i, u_j} S_{0,1}^-(u)\}.$$

Minimum of $M_2'^+, M_2''^+, M_2'''^+$ and $M_2''''^+$ is the best value for $m_1^+(v)$. So:

$$m_1^+(v) = \text{Min}\{M_2'^+, M_2''^+, M_2'''^+, M_2''''^+\}.$$

Calculating $m_0^-(v)$ when v is not a pre-pendent

In this case, v is white and none of the child of v and $ch(v)$ are not black. If v has a child like u that is a none empty block node, then vertices of u can not dominated by any vertices because v or $ch(u)$ can only dominate u . So, $m_0^-(v) = \infty$ otherwise we have:

$$m_1^-(v) = \sum_{u \in ch(v)} S_{1,2}^-(u).$$

Calculating $m_1^-[v]$ when v is not a pre-pendent

In this case, v is white and exactly one of the child of v or $ch(v)$ are black.

- The node v has at least two children like u_1 and u_2 that are none empty block node. So vertices of u_1 or vertices of u_2 can not dominated and $m_1^-(v) = \infty$.
- The node v has only one none empty child like u_1 and the other children are empty. So two cases appear:
 1. One of the vertex of block node u_1 is black and all of children of u_1 are white and are dominated at most once.
 2. All vertices of block node u_1 are white, exactly one child v_i of u_1 is black, the other are white. Moreover, v_i is dominated once or twice and white siblings of v_i are dominated at most once.

And we have:

$$m_1^-(v) = \text{Min}\{1 + S_{0,1}^-(u_1) + \sum_{u \in ch(v), u \neq u_1} S_{1,2}^-(u), S_{0,1}^+(u) + \sum_{u \in ch(v), u \neq u_1} S_{1,2}^-(u)\}$$

- All children of v are empty. So, among all children of v , there is exactly one node u_i such that u_i has only a black child the other is white. In other child $u \neq u_i$, all nodes of $ch(u)$ should be white and we have:

$$m_1^-(v) = \text{Min}_{u_i \in ch(v)} \{S_{0,1}^+(u_i) + \sum_{u \in ch(v), u \neq u_i} S_{1,2}^-(u)\}$$

All children of v are empty. So, from exactly one child u_i of v , one of the children is black and the others are white. In other child $u \neq u_i$, all nodes of $ch(u)$ should be white and we have:

$$m_1^-(v) = \text{Min}_{u_i \in ch(v)} \{S_{0,1}^+(u_i) + \sum_{u \in ch(v), u \neq u_i} S_{1,2}^-(u)\}$$

Calculating $m_2^-[v]$ when v is not a pre-pendent

In the last cases, we consider v is white and exactly two children of v or $ch(v)$ are black.

- The node v has more than two none empty children, $m_2^-[v] = \infty$.
- The node v has only two none empty children like u_1 and u_2 and the other children are empty. So, for u_1, u_2 and the other vertices one of the following cases appears:
 1. One of the vertex of block nodes u_1 and u_2 are black, all of children of them are white and are dominated at most once.
 2. One of the vertex of block node u_1 is black, all of children of it are white and are dominated at most once. Moreover, all of the vertices of block node u_2 are white, one of its child is black, the others are white and dominated at most once. (u_1 and u_2 can replace.)
 3. All of the vertices of block node u_1, u_2 are white, one of its child is black, the others are white and dominated at most once.

So we have:

$$\begin{aligned}
 m_2^-(v) = \text{Min}\{ & 2 + S_{0,1}^-(u_1) + S_{0,1}^-(u_2) + \sum_{u \in ch(v), u \neq u_1, u_2} S_{1,2}^-, \\
 & 1 + S_{0,1}^+(u_1) + S_{0,1}^-(u_2) + \sum_{u \in ch(v), u \neq u_1, u_2} S_{1,2}^-, \\
 & 1 + S_{0,1}^+(u_2) + S_{0,1}^-(u_1) + \sum_{u \in ch(v), u \neq u_1, u_2} S_{1,2}^-, \\
 & S_{0,1}^+(u_1) + S_{0,1}^+(u_2) + \sum_{u \in ch(v), u \neq u_1, u_2} S_{1,2}^- \}
 \end{aligned}$$

- The node v has only one none empty child u_1 . So, one of the following cases occurs for u_1 :
 1. Two children of u_1 are black and the others are white and not dominated at most once.
 2. One vertex of block node u_1 and one of its child are black and the other children of u_1 are white and not dominated.
 3. One vertex of block node u_1 is black and all of children of u_1 are white and are dominated at most once.
 4. All vertices of block node u_1 are white, one of its child are black and the other children of u_1 are white and not dominated.

For other child u of $ch(v)$, there are exactly one node u_i such that one child of it is black and the others are white. Obviously, all child of other siblings u_i are white. So we have:

$$\begin{aligned}
 m_2^-(v) = \text{Min}\{ & S^*(u_1) + \sum_{u \in ch(v), u \neq u_1} S_{1,2}^-(u_1), \\
 & 1 + S_0^+(u_1) + \sum_{u \in ch(v), u \neq u_1} S_{1,2}^-(u_1), \\
 & 1 + S_{0,1}^-(u_1) + \text{Min}_{u_i \in ch(v), u_i \neq u_1} \{ S_{0,1}^+(u_i) + \sum_{u \in ch(v), u \neq u_i, u_1} S_{1,2}^-(u) \}, \\
 & S_{0,1}^+(u_1) + \text{Min}_{u_i \in ch(v), u_i \neq u_1} \{ S_{0,1}^+(u_i) + \sum_{u \in ch(v), u \neq u_i, u_1} S_{1,2}^-(u) \} \}.
 \end{aligned}$$

- All children of v are empty. So, from exactly two children u_i, u_j of v , one of the children is black and the others are white. In other child $u \neq u_i, u_j$, all nodes of $ch(u)$ should be white and we have:

$$m_2^-(v) = \text{Min}_{u_i, u_j \in ch(v)} \{ S_{0,1}^+(u_i) + S_{0,1}^+(u_j) + \sum_{u \in ch(v), u \neq u_i, u_j} S_{1,2}^-(u) \}$$

2.5. Final state:

Let r be the root of refined cut tree T , r can be correspond to a cut vertex of G or a block of it. Depend on type of r one of the following cases appear:

1. The root r of T is a cut vertex of G .

Since for $i = 0, 1, 2$ we compute $m_i^+[v]$ and $m_i^-[v]$ on a node of T that its corresponding vertex in G is a cut vertex, so we must choose best set among computed set of root r . Note that r should be black or white and should be dominated by one or two vertices. It means that:

$$M = \text{Min}\{m_1^+(r), m_2^+(r), m_1^-(r), m_2^-(r)\}.$$

2. The root r of T is corresponding to a block of G .

Since, we computed $m^+[v], m_0^-[v], m_1^-[v]$ and $m_2^-[v]$ for all vertices $v \in ch(r)$. Based on the number of vertices in block r and the number of its child, one of the following cases appear:

- (a) $|r| = 0$, so we have:

$$M = \text{Min}\{S_{1,2}^-(r), S_{0,1}^+(r), S^*(r)\}.$$

- (b) $|r| > 0$, so we have::

$$M = \text{Min}\{1 + S_0^+(r), 1 + S_{0,1}^-(r), S_{0,1}^+(r), S_{1,2}^-(r), S^*(r)\}.$$

Theorem 2.1. *The value M computed by Algorithm 1 for the block graph G is size of the smallest total $[1, 2]$ -set of G and is computed in linear-time.*

Proof. The number of nodes in the refined cut tree T corresponding to block graph G is linear based on order of G , i.e. n . It is obvious that the algorithm traverses T once and is computed in linear-time $O(n)$. □

Algorithm 1 Total $[1, 2]$ -Dominating Set

Input: A refined cut tree T of block graph G .

- 1: **procedure** INITIALIZING STEP
- 2: **procedure** UPDATING STEP: \triangleright Depending on the type of non pre-pendant nodes in the post order traversal of T , one of the following procedures is selected:
- 3: **procedure** UPDATING STEP FOR BLOCK NODES OF T :
- 4: Calculating $S_0^-(u), S_0^+(u), S_{0,1}^-(u), S_{1,2}^-(u), S_0^+(u), S_{0,1}^+(u)$ and $S^*(u)$.
- 5: **procedure** UPDATING STEP FOR CUT VERTEX OF T
- 6: Calculating $m_0^+(v), m_1^+(v), m_0^-(v), m_1^-[v]$ and $m_2^-[v]$ when v is not a pre-pendent.
- 7: **procedure** FINAL STATE
- 8: **if** the root r of T is a cut vertex of G **then** $M = \text{Min}\{m_1^+(r), m_2^+(r), m_1^-(r), m_2^-(r)\}$.
- 9: **if** the root r is a block and $|r| = 0$ **then** $M = \text{Min}\{S_{1,2}^-(r), S_{0,1}^+(r), S^*(r)\}$.
- 10: **if** the root r is a block and $|r| > 0$ **then**

$$M = \text{Min}\{1 + S_0^+(r), 1 + S_{0,1}^-(r), S_{0,1}^+(r), S_{1,2}^-(r), S^*(r)\}.$$

Output: Size of minimum total $[1, 2]$ -set of G .

References

- [1] A. V. Aho, J. E. Hopcroft, The design and analysis of computer algorithms, Pearson Education India (1974).
- [2] A. Bishnu, K. Dutta, A. Gosh, S. Pual, $[1, j]$ -set problem in graphs, Discrete Mathematics 339 (2016), 2215-2525.

- [3] M. Chellali, O. Favaron, T. W. Haynes, S. T. Hedetniemi, A. McRae, Independent $[1, k]$ -sets in graphs, *Australasian Journal of Combinatorics* 59, 1 (2014), 144-156.
- [4] M. Chellali, T. W. Haynes, S. T. Hedetniemi, A. McRae, $[1, 2]$ -sets in graphs, *Discrete Applied Mathematics* 161, 18 (2013), 2885-2893.
- [5] O. Etesami, N. Ghareghani, M. Habib, M. R. Hooshmandasl, R. Naserasr, P. Sharifani, When an optimal dominating set with given constraints exists, *Theoretical Computer Science*, 780 (2019), 54-65.
- [6] A. K. Goharshady, M. R. Hooshmandasl, M. Alambardar Meybodi, $[1, 2]$ -sets and $[1, 2]$ -total sets in trees with algorithms, *Discrete Applied Mathematics*, 198 (2016), 136-146.
- [7] T. W. Haynes, S. T. Hedetniemi, P. J. Slater, *Fundamentals of domination in graphs*, CRC Press, 1998.
- [8] T. W. Haynes, S. T. Hedetniemi, P. J. Slater, *Domination in graphs: advanced topics*, Marcel Dekker, 1998.
- [9] S. T. Hedetniemi, R. Laskar, *Topics on domination*, Elsevier, 1991.
- [10] D. B. West, *Introduction to graph theory*, 2ed Edition. Prentice hall Upper Saddle River, 2001.
- [11] X. Yang, B. Wu, $[1, 2]$ -domination in graphs, *Discrete Applied Mathematics*, 175 (2014), 79-86.

Please cite this article using:

Pouyeh Sharifani, Mohammadreza Hooshmandasl, Saeid Alikhani, A linear-time algorithm to compute total $[1, 2]$ -domination number of block graphs, *AUT J. Math. Comput.*, 1(2) (2020) 263-270
DOI: 10.22060/ajmc.2020.18444.1035

