



## A note on the Yamabe problem of Randers metrics

Bin Chen<sup>\*a</sup>, Siwei Liu<sup>a</sup>

<sup>a</sup>*School of Mathematical Sciences, Tongji University, Shanghai, China, 200092*

**ABSTRACT:** The classical Yamabe problem in Riemannian geometry states that every conformal class contains a metric with constant scalar curvature. In Finsler geometry, the C-convexity is needed in general. In this paper, we study the strong C-convexity of Randers metrics, and provide a result on the Yamabe problem for the metrics of Randers type.

### Review History:

Received:23 June 2021

Accepted:24 July 2021

Available Online:01 September 2021

### Keywords:

Randers metrics

C-convex

Yamabe problem

### AMS Subject Classification (2010):

53C60; 58B20

*(Dedicated to Professor Zhongmin Shen for his warm friendship and research collaboration)*

### 1. Introduction

H. Yamabe attempted to seek Riemannian metrics with constant scalar curvature in a conformal class in [11]. A bundle of works of N. Trudinger [10], T. Aubin [1] and R. Schoen [9] gives an affirmative answer to the Yamabe Problem, which is a milestone in Riemannian geometry. In Finsler realm, X. Cheng and M. Yuan [4] studied the Yamabe problem for the scalar curvature defined by H. Akbar-Zadeh, and obtained a negative answer for Randers metrics. In the view of calculus of variations, L. Zhao and the first author defined a Finsler scalar curvature  $\text{Scal}(x)$  and proved that a Finsler metric with constant scalar curvature is a critical point of the total scalar curvature functional

$$\mathcal{S}(F) = \frac{1}{\text{Vol}(M)^{1-\frac{2}{n}}} \int_M \text{Scal}(x) d\mu_F$$

in its conformal class([5]). The Yamabe invariant is defined as  $Y(M, F) = \inf_u \mathcal{S}(e^{u(x)}F)$ . In order to have a lower bound of  $\mathcal{S}$  in the conformal class  $[F]$  of the metric  $F$ , the condition *C-convex* is introduced in [5] which is conformally invariant. By introducing another conformal invariant  $C(M, F)$ , L. Zhao and the first author partially solved the Yamabe problem in Finsler geometry.

**Theorem 1.1 ([5]).** *Let  $(M^n, F)$  be a compact C-convex Finsler manifold with  $n \geq 3$ . If  $Y(M, F)C(M, F) < Y(\mathbb{S}^n)$ , then there exists a metric  $\bar{F}$  conformal to  $F$  such that  $\text{Scal}_{\bar{F}}(x) = Y(M, \bar{F})$ .*

<sup>\*</sup>Corresponding author.

E-mail addresses: chenbin@tongji.edu.cn, carrot98@163.com

Supported by the National Natural Science Foundation of China (No. 11871126)

In [6], L. Zhao and the first author introduce the notion of *strongly C-convexity* (see (2.1) in §2) which is a bit more stronger than C-convexity. In this paper, we shall study the strong C-convexity of Randers metrics and obtain the following result.

**Theorem 1.2.** *A Randers metric  $\alpha + \beta$  is strongly C-convex if and only if*

$$\|\beta\|_\alpha < B_n := \left(1 - \left(\frac{8n^3 - 16n^2 + 8n + 2}{n^4 + 3n^3 - 9n^2 + 7n}\right)^2\right)^{1/2}.$$

where the dimension  $n \geq 3$ .

As a conclusion, we can give a statement to the Yamabe Problem of Randers metrics.

**Corollary 1.3.** *Let  $F = \alpha + \beta$  be a Randers metric on a compact manifold  $M^n$  with  $n \geq 3$ . If  $\|\beta\|_\alpha < B_n$  and  $Y(M, F)C(M, F) < Y(S^n)$ , then there exists a metric  $\bar{F}$  conformal to  $F$  such that  $\text{Scal}_{\bar{F}}(x) = Y(M, F)$ .*

The contents of this paper are arranged as follows. In §2, we give a brief review of Finsler metrics and give the precise definition of strongly C-convex. In §3, we study the strongly C-convexity of Randers metrics. Throughout this paper, we always assume that the dimension  $n \geq 3$ .

## 2. Finsler metrics

Let  $M$  be an  $n$ -dimensional differentiable manifold with  $n \geq 3$ . The points in the tangent bundle  $TM$  are denoted by  $(x, y)$ , where  $x \in M$  and  $y \in T_x M$ . Let  $(x^i; y^i)$  be the local coordinates of  $TM$  with  $y = y^i \partial / \partial x^i$ .

Let  $F : TM \rightarrow [0, +\infty)$  be a Finsler metric on  $M$ . The fundamental form of  $F$  is

$$g = g_{ik}(x, y) dx^i \otimes dx^k, \quad g_{ik} := \left[ \frac{1}{2} F^2 \right]_{y^i y^k}.$$

Here and from now on, the lower index  $x^i, y^i$  always means partial derivatives, such as  $F_{y^i} := \frac{\partial F}{\partial y^i}$ ,  $F_{x^i} := \frac{\partial F}{\partial x^i}$ ,  $[F^2]_{y^i y^k} := \frac{\partial^2 F^2}{\partial y^i \partial y^k}$ , and etc.

The canonical projection  $\pi : TM \setminus \{0\} \rightarrow M$  gives rise to a covector bundle  $\pi^* T^* M$ , on which there exists the Hilbert form  $\omega = \ell_i dx^i$  where  $\ell_i = F_{y^i}$ , whose dual is the distinguished section of  $\pi^* T^* M$

$$\ell = \ell^i \frac{\partial}{\partial x^i}, \quad \text{with } \ell^i := \frac{y^i}{F}.$$

The Cartan tensor (Cartan torsion) and the Cartan form are respectively

$$A = A_{ijk} dx^i \otimes dx^j \otimes dx^k, \quad A_{ijk} := \frac{F}{4} [F^2]_{y^i y^j y^k},$$

$$I = I_i dx^i, \quad I_i := A_{ijk} g^{jk}, \quad (g^{jk}) = (g_{jk})^{-1}.$$

The spray coefficients are given as

$$G^i = \frac{1}{4} g^{il} \{ [F^2]_{x^k y^l y^k} - [F^2]_{x^l} \}$$

which determine the Berwald connection coefficients in the following way

$$\Gamma_{jk}^i = G_{y^j y^k}^i.$$

The flag curvature tensor (Riemann curvature tensor) is given by

$$R^i_k = 2G_{x^k}^i - G_{x^j y^k}^i y^j + 2G^j G_{y^j y^k}^i - G_{y^j}^i G_{y^k}^j,$$

while the Ricci curvature is defined as the trace

$$\text{Ric}(x, y) := \frac{1}{F^2} R^i_i.$$

The most important non-Riemannian curvature in Finsler geometry is the Landsberg curvature, which is defined as the derivative of the Cartan torsion

$$L_{ijk} := A_{ijk:m} \ell^m$$

where “:” is the horizontal covariant derivative with respect to the Berwald connection. The mean Landsberg tensor is

$$J = J_k dx^k, \quad J_k := g^{ij} L_{ijk}.$$

On the punctured bundle  $TM \setminus \{0\}$ , there is the Sasaki type metric  $g_{ik} dx^i \otimes dx^k + g_{ik} \frac{\delta y^i}{F} \otimes \frac{\delta y^k}{F}$ , which induces a Riemannian metric on the projective sphere bundle  $SM$

$$\hat{g} = g_{ik} dx^i \otimes dx^k + F[F]_{y^i y^k} \frac{\delta y^i}{F} \otimes \frac{\delta y^k}{F}.$$

Hence the volume form of  $SM$  can be expressed as [3, 7]

$$d\mu_{SM} = \Omega d\eta \wedge dx, \quad \Omega := \det \left( \frac{g_{ik}}{F} \right)$$

where

$$d\eta := \sum (-1)^{i-1} y^i dy^1 \wedge \cdots \wedge \widehat{dy^i} \wedge \cdots \wedge dy^n, \quad dx := dx^1 \wedge \cdots \wedge dx^n.$$

The volume form of  $M$  induced by  $SM$  can be defined by

$$d\mu_F = \sigma_F(x) dx, \quad \sigma_F(x) := \frac{1}{\omega_{n-1}} \int_{S_x M} \Omega d\eta,$$

where  $\omega_{n-1}$  is the volume of the  $(n - 1)$ -dimensional standard sphere.

By integrating along the fibre, the scalar curvature can be defined as

$$\text{Scal}(x) = \frac{n \int_{S_x M} Ric \cdot \Omega d\eta}{\int_{S_x M} \Omega d\eta} + \frac{2n}{n-2} \frac{\int_{S_x M} g^{ij} J_{i;j} \cdot \Omega d\eta}{\int_{S_x M} \Omega d\eta}$$

as the dimension  $n \geq 3$ . In order to obtain the existence of Finsler metrics with constant scalar curvature  $\text{Scal}(x)$ , the concept of C-convexity is introduced in [5]. Precisely, a Finsler metric is *strongly C-convex* if the tensor

$$\mathfrak{C}^{ij} = g^{ij} - \frac{n}{(n-1)(n-2)} (\ell^i I^j + \ell^j I^i + A_s^{ir} A_r^{js}) \tag{2.1}$$

is positive definite, while *C-convexity* means the positivity of the tensor

$$c^{ij} = \frac{1}{\int_{S_x M} \Omega d\eta} \int_{S_x M} \mathfrak{C}^{ij} \cdot \Omega d\eta.$$

We shall point out that the C-convexity does not make sense as  $n = 2$ .

One can find that a metric is strongly C-convex if its Cartan torsion is sufficiently small. In the next section, we shall study the strongly C-convexity of Randers metrics and obtain Theorem 1.2.

### 3. Strongly C-convexity of Randers Metrics

Let  $F = \alpha + \beta$  be a Randers metric where  $\alpha = \sqrt{a_{ij} y^i y^j}$  and  $\beta = b_i y^i$  with  $b = \sqrt{a^{ij} b_i b_j} < 1$  and  $(a^{ij}) = (a_{kl})^{-1}$ . In this section, we shall investigate the positivity of  $(\mathfrak{C}^{ij})$  of  $F = \alpha + \beta$ .

It is well-known that the Cartan tensor of a Randers metric is reducible

$$A_{ijk} = \frac{1}{n+1} \{I_i h_{ik} + I_j h_{ik} + I_k h_{ij}\}$$

where  $h_{ij} := FF_{y^i y^j} = g_{ij} - \ell_i \ell_j$  is the angular tensor, and the Cartan form is

$$I_i = \frac{n+1}{2} \left( b_i - \frac{\beta}{\alpha} \alpha_{y^i} \right).$$

Being aware of  $h_s^r h_r^s = n - 1$  and  $h_s^i h^{js} = g^{ij} - \ell^i \ell^j$ , one can get

$$A_s^{ir} A_r^{js} = \frac{1}{(n+1)^2} \{2\|I\|^2 g^{ij} + (n+5)I^i I^j - 2\|I\|^2 \ell^i \ell^j\}$$

where

$$\|I\|^2 = I_i I_j g^{ij} = \frac{(n+1)^2}{4} \cdot \frac{b^2 - (\beta/\alpha)^2}{1 + (\beta/\alpha)} \leq \frac{(n+1)^2}{2} (1 - \sqrt{1 - b^2}), \tag{3.1}$$

which can be found in [8]. Thus, we reach

$$\begin{aligned} \mathfrak{C}^{ij} &= \left( 1 - 2\|I\|^2 \frac{n}{(n+1)^2(n-1)(n-2)} \right) g^{ij} \\ &\quad - \frac{n}{(n-1)(n-2)} \left( \ell^i I^j + \ell^j I^i + \frac{n+5}{(n+1)^2} I^i I^j - \frac{2\|I\|^2}{(n+1)^2} \ell^i \ell^j \right). \end{aligned}$$

For investigating the positivity, one can apply the continuity method. Let us consider the family  $F_t = \alpha + t\beta$  where  $t \in [0, 1]$ . It is clear that  $(\mathfrak{C}^{ij})$  of  $F_0$  is  $(a^{ij})$  which is positive definite. Hence, once we obtain the invertibility of  $(\mathfrak{C}^{ij})$  for every  $F_t$ , we shall have the positivity of  $(\mathfrak{C}^{ij})$  for every  $F_t$ . Thus we shall calculate the determinant  $\det(\mathfrak{C}^{ij})$  by applying the following lemma.

**Lemma 3.1 ([2]).** *Let  $H = (H^{ij})$  be a symmetric  $n \times n$  matrix and  $V = (V^i)$  be an  $n$ -vector. Put  $G^{ij} = H^{ij} + \delta V^i V^j$  where  $\delta$  is a complex number. Assuming that  $H$  is invertible with  $H^{-1} = (H_{ij})$ , it holds*

$$\det(G^{ij}) = (1 + \delta v) \det(H^{ij}),$$

where  $v = V_i V^i$  and  $V_i = H_{ij} V^j$ . Moreover, if  $1 + \delta v \neq 0$ ,  $G$  is invertible and the inverse  $G^{-1} = (G_{ij})$  is given by

$$G_{ij} = H_{ij} - \frac{\delta V_i V_j}{1 + \delta v}.$$

In order to apply the above lemma, we rewrite  $\mathfrak{C}^{ij}$  in the following form

$$\mathfrak{C}^{ij} = \rho_0 g^{ij} + \rho_1 \ell^i \ell^j - \rho_2 \left( I^i + \frac{(n+1)^2}{n+5} \ell^i \right) \left( I^j + \frac{(n+1)^2}{n+5} \ell^j \right) \tag{3.2}$$

where

$$\begin{aligned} \rho_0 &= 1 - 2\|I\|^2 \frac{n}{(n+1)^2(n-1)(n-2)}, \\ \rho_1 &= \frac{n}{(n-1)(n-2)} \left( \frac{(n+1)^2}{n+5} + \frac{2\|I\|^2}{(n+1)^2} \right), \\ \rho_2 &= \frac{n(n+5)}{(n+1)^2(n-1)(n-2)}. \end{aligned}$$

**Lemma 3.2.** *The coefficient  $\rho_0$  is positive when  $n \geq 4$ . In dimension  $n = 3$ ,  $\rho_0 > 0$  if and only if  $b^2 < \frac{8}{9}$ .*

**Proof.** By (3.1), we have  $\|I\|^2 < \frac{(n+1)^2}{2}$ , thus for  $n \geq 4$  we have

$$\rho_0 = 1 - 2\|I\|^2 \frac{n}{(n+1)^2(n-1)(n-2)} > 1 - \frac{n}{(n-1)(n-2)} > 0.$$

For  $n = 3$ , we use the estimate  $\|I\|^2 \leq \frac{(n+1)^2}{2} (1 - \sqrt{1 - b^2})$  where the equality can be achieved. Thus  $\rho_0 > 0$  if and only if

$$\min_y \rho_0 = 1 - \frac{3}{16} \max_y \|I\|^2 = 1 - \frac{3}{2} (1 - \sqrt{1 - b^2}) > 0$$

which implies  $b^2 < \frac{8}{9}$ . □

In the remaining part of this section, we shall assume  $\rho_0 > 0$ . Therefore, by putting

$$\tilde{H}^{ij} = \rho_0 g^{ij} + \rho_1 \ell^i \ell^j = \rho_0 \left( g^{ij} + \frac{\rho_1}{\rho_0} \ell^i \ell^j \right),$$

and according to Lemma 3.1, we have

$$\det(\tilde{H}^{ij}) = (\rho_0)^n \det(g^{ij}) \left( 1 + \frac{\rho_1}{\rho_0} \right).$$

One can easily find that  $1 + \frac{\rho_2}{\rho_0} > 0$ . Thus  $(\tilde{H}^{ij})$  is invertible with the inverse

$$\tilde{H}_{ij} = \frac{1}{\rho_0} g_{ij} - \frac{\rho_1}{(\rho_0)^2} \frac{\ell_i \ell_j}{1 + \frac{\rho_1}{\rho_0}}.$$

Now, applying Lemma 3.1 to

$$\mathfrak{C}^{ij} = \tilde{H}^{ij} - \rho_2 \left( I^i + \frac{(n+1)^2}{n+5} \ell^i \right) \left( I^j + \frac{(n+1)^2}{n+5} \ell^j \right),$$

we obtain

$$\begin{aligned} \det(\mathfrak{C}^{ij}) &= \det(\tilde{H}^{ij}) \left[ 1 - \rho_2 \tilde{H}_{ij} \left( I^i + \frac{(n+1)^2}{n+5} \ell^i \right) \left( I^j + \frac{(n+1)^2}{n+5} \ell^j \right) \right] \\ &= (\rho_0)^n \det(g^{ij}) \left[ 1 - \frac{\rho_2}{\rho_0} \|I\|^2 - \frac{(n+1)^4}{(n+5)^2} \frac{\rho_2}{\rho_0 + \rho_1} \right] \\ &= (\rho_0)^n \det(g^{ij}) \left[ 1 - \frac{n(n+1)^2}{2n^3 + 4n^2 - 12n + 10} \right. \\ &\quad \left. - \frac{n(n+5)\|I\|^2}{(n+1)^2(n-1)(n-2) - 2n\|I\|^2} \right]. \end{aligned}$$

**Theorem 3.3.** *A Randers metric  $F$  is strongly C-convex if and only if*

$$b < B_n := \left( 1 - \left( \frac{8n^3 - 16n^2 + 8n + 2}{n^4 + 3n^3 - 9n^2 + 7n} \right)^2 \right)^{1/2}$$

where  $b = \|\beta\|_\alpha$ .

**Proof.** By the decomposition (3.2) and  $n \geq 3$ , the positivity of  $(\mathfrak{C}^{ij})$  shall imply  $\rho_0 > 0$ . In fact, since  $n \geq 3$ , by picking a covector  $(V_i)$  such that

$$\ell^i V_i = 0, \quad \left( I^i + \frac{(n+1)^2}{n+5} \ell^i \right) V_i = 0,$$

we have  $\mathfrak{C}^{ij} V_i V_j = \rho_0 g^{ij} V_i V_j$ . Thus the positivity of  $(\mathfrak{C}^{ij})$  does imply  $\rho_0 > 0$ . Hence, if  $F$  is strongly C-convex, we have  $\rho_0 > 0$  and  $\det(\mathfrak{C}^{ij}) > 0$ . Therefore, one shall get

$$1 - \frac{n(n+1)^2}{2n^3 + 4n^2 - 12n + 10} - \frac{n(n+5)\|I\|^2}{(n+1)^2(n-1)(n-2) - 2n\|I\|^2} > 0. \tag{3.3}$$

The inequality (3.3) is equivalent to

$$\|I\|^2 < \frac{(n+1)^2(n-1)(n-2)(n(n-2)-1)}{2n(n^3+3n^2-9n+7)}.$$

Since  $\max_y \|I\|^2 = \frac{(n+1)^2}{2}(1 - \sqrt{1-b^2})$ , the above inequality holds if and only if

$$\frac{(n+1)^2}{2}(1 - \sqrt{1-b^2}) < \frac{(n+1)^2(n-1)(n-2)(n(n-2)-1)}{2n(n^3+3n^2-9n+7)},$$

from which we can get

$$b^2 < 1 - \left( \frac{8n^3 - 16n^2 + 8n + 2}{n^4 + 3n^3 - 9n^2 + 7n} \right)^2. \tag{3.4}$$

Conversely, let us assume that  $\beta$  satisfies (3.4). A simple calculation shows that  $b^2 < \frac{8}{9}$  as  $n = 3$ . Hence,  $\rho_0$  is positive according to Lemma 3.2. Since (3.15) is equivalent to (3.18), we have  $\det(\mathfrak{C}^{ij}) > 0$  in this case. Now, put  $F_t = \alpha + t\beta$  for  $t \in [0, 1]$ , and denote  $\mathfrak{C}^{ij}$  of  $F_t$  by  $\mathfrak{C}^{ij}(t)$ . It is clear that  $t\beta$  also satisfies (3.18) since  $\|t\beta\|_\alpha \leq \|\beta\|_\alpha$ . Hence, for every  $t \in [0, 1]$  we have  $\det(\mathfrak{C}^{ij}(t)) > 0$  and thus the eigenvalues of  $\mathfrak{C}^{ij}(t)$  are nonzero. Note that the eigenvalues of  $\mathfrak{C}^{ij}(t)$  depend continuously on  $t$ , and  $\mathfrak{C}^{ij}(0) = \alpha^{ij}$  is positive definite. As  $t$  changes from 0 to 1, none of these eigenvalues can become negative. Thus  $\mathfrak{C}^{ij}$  is positive definite. □

**Remark.** In order to have an intuition, we list below the decimal values of several  $B_n$ 's.

	$n = 3$	$n = 4$	$n = 5$	$n = 10$	$n = 100$	$n = 1000$
$B_n$	0.2773	0.4869	0.6098	0.8464	0.9971	0.9999

As  $n$  grows, the condition on  $\beta$  becomes weaker. While it is critical as the dimension is low.

## References

- [1] T. Aubin, Equations differentielles non lineaires et probleme de Yamabe concernant la courbure scalaire, *J. Math. Pures Appl.* 55 (1976), 269-296.
- [2] D. Bao, S. -S. Chern and Z. Shen, An introduction to Riemann-Finsler geometry, *Grad. Texts in Math.* 200, Springer, New York, 2000.
- [3] B. Chen and Y. Shen, On a class of critical Riemann-Finsler metrics, *Publ. Math. Debrecen*, 72/3-4 (2008), 451-468.
- [4] X. Cheng and M. Yuan, On Randers metrics of isotropic scalar curvature, *Publ. Math. Debrecen*, 84/1-2 (2014), 63-74.
- [5] B. Chen and L. Zhao, On a Yamabe type problem in Finsler geometry, *Canad. Math. Bull.* 60(2) (2017), 253-268.
- [6] B. Chen and L. Zhao, Finsler conformal changes preserving the modified Ricci curvature, *Tohoku Math. J.*, 73 (2021), 39-48.
- [7] Q. He and Y. Shen, Some results on harmonic maps for Finsler manifolds, *Inter. Jour. of Math.*, Vol.16, No.9(2005), 1017-1031.
- [8] M. Ji and Z. Shen, On strongly convex graphs in Minkowski geometry, *Canad. Math. Bull.* 45(2) (2002), 232-246.
- [9] R. Schoen, Conformal deformation of a Riemannian metric to constant scalar curvature, *J. Diff. Geom.* 20 (1984), 479-495.
- [10] N. Trudinger, Remarks concerning the conformal deformation of Riemannian structures on compact manifolds, *Ann. Scuola Norm. Sup. Pisa* 22 (1968), 265-274.
- [11] H. Yamabe, On the deformation of Riemannian structures on compact manifolds, *Osaka Math. Jour.*, 12 (1960), 21-37.

Please cite this article using:

Bin Chen, Siwei Liu, A note on the Yamabe problem of Randers metrics, *AUT J. Math. Comput.*, 2(2) (2021) 165-170  
DOI: 10.22060/ajmc.2021.20199.1056

