



Original Article

Bergman and Dirichlet spaces in the unit ball and symmetric lifting operator

Mostafa Hassanlou<sup>a</sup>, Ebrahim Abbasi<sup>\*b</sup>

<sup>a</sup>Engineering Faculty of Khoy, Urmia University of Technology, Urmia, Iran

<sup>b</sup>Department of Mathematics, Mahabad Branch, Islamic Azad University, Mahabad, Iran

**ABSTRACT:** Let  $\mathbb{B}_n$  be the open unit ball in  $\mathbb{C}^n$  and  $\mathbb{B}_n^2 = \mathbb{B}_n \times \mathbb{B}_n$ . The symmetric lifting operator which lifts analytic functions from  $H(\mathbb{B}_n)$  to  $H(\mathbb{B}_n^2)$  is defined as follow

$$L(f)(z, w) = \frac{f(z) - f(w)}{z - w}.$$

In this paper we investigate the action of symmetric lifting operator on the Bergman space in the unit ball. Also, we state a characterization for Dirichlet space and consider symmetric lifting operator on the Dirichlet space in the unit ball.

**Review History:**

Received:16 September 2022  
Revised:09 December 2022  
Accepted:09 December 2022  
Available Online:20 March 2023

**Keywords:**

Symmetric lifting operator  
Bergman space  
Dirichlet space  
Pseudo-hyperbolic metric

**AMS Subject Classification (2010):**

16W25; 46L57; 47B47

**1. Introduction**

For  $z = (z_1, \dots, z_n)$  and  $w = (w_1, \dots, w_n)$  in  $\mathbb{C}^n$ , we define  $\langle z, w \rangle = z_1 \overline{w_1} + \dots + z_n \overline{w_n}$ , where  $\overline{w_k}$  is the complex conjugate of  $w_k$ . We also write  $|z| = \sqrt{\langle z, z \rangle} = \sqrt{|z_1|^2 + \dots + |z_n|^2}$ . Let  $\mathbb{B}_n$  denote the open unit ball of  $\mathbb{C}^n$ , that is

$$\mathbb{B}_n = \{z \in \mathbb{C}^n : |z| < 1\}.$$

Thus for any  $a \in \mathbb{B}_n - \{0\}$ , we define

$$\varphi_a(z) = \frac{a - P_a(z) - s_a Q_a(z)}{1 - \langle z, a \rangle} \quad z \in \mathbb{B}_n,$$

where  $s_a = \sqrt{1 - |a|^2}$ ,  $P_a$  is the orthogonal projection from  $\mathbb{C}^n$  onto the subspace  $[a]$  generated by  $a$ , and  $Q_a$  is the orthogonal projection from  $\mathbb{C}^n$  onto  $\mathbb{C}^n - [a]$ . When  $a = 0$ , write  $\varphi_a(z) = -z$ . These functions are called

\*Corresponding author.

E-mail addresses: mst8084@gmail.com, ebrahimabbasi81@gmail.com



involutions.

The hyperbolic metric(Bergman metric) is defined by

$$\beta(z, w) = \frac{1}{2} \log \frac{1 + |\varphi_z(w)|}{1 - |\varphi_z(w)|}, \quad z, w \in \mathbb{B}_n.$$

For any  $z \in \mathbb{B}_n$  and  $r > 0$ , we denote Bergman metric ball at  $z$  by  $D(z, r)$ . That is

$$D(z, r) = \{w \in \mathbb{B}_n : \beta(z, w) < r\}.$$

Also, pseudo-hyperbolic metric defined as  $\rho(z, w) = |\varphi_z(w)|$ .

For  $\alpha > -1$  let

$$dv_\alpha(z) = c_\alpha(1 - |z|^2)^\alpha dv(z),$$

where  $dv$  is the Lebesgue volume measure on  $\mathbb{B}_n$  and  $c_\alpha$  is a positive constant with  $v_\alpha(\mathbb{B}_n) = 1$ . For  $0 < p < \infty$  and  $\alpha > -1$ , the weighted Bergman space  $A_\alpha^p(\mathbb{B}_n)$  consists of all holomorphic functions in  $L^p(\mathbb{B}_n, dv_\alpha)$ , that is

$$A_\alpha^p(\mathbb{B}_n) = \left\{ f \in H(\mathbb{B}_n) : \|f\|_{\alpha,p}^p = \int_{\mathbb{B}_n} |f(z)|^p dv_\alpha(z) < \infty \right\}.$$

More information about Bergman spaces can be found in [1, 4, 7, 9]. Let  $\mathbb{B}_n^2 = \mathbb{B}_n \times \mathbb{B}_n$  be the open subset of  $\mathbb{C}_n^2 = \mathbb{C}_n \times \mathbb{C}_n$  which is

$$\mathbb{B}_n^2 = \{(z, w) \in \mathbb{C}_n^2 : |z| < 1, |w| < 1\}.$$

The Bergman space  $A_\alpha^p(\mathbb{B}_n^2)$  over the  $\mathbb{B}_n^2$  is the space of all holomorphic functions on  $\mathbb{B}_n^2$  such that

$$\int_{\mathbb{B}_n} \int_{\mathbb{B}_n} |f(z, w)|^p dv_\alpha(z) dv_\alpha(w) < \infty.$$

For  $f \in \mathbb{B}_n$ , some notations which will be used in this section are:

$$Rf(z) = \sum_{k=1}^n z_k \frac{\partial f}{\partial z_k}(z).$$

$Rf$  is called the radial derivative of  $f$ . The (holomorphic) gradient of  $f$  at  $z$  is

$$|\nabla f(z)| = \left| \left( \frac{\partial f}{\partial z_1}(z), \dots, \frac{\partial f}{\partial z_n}(z) \right) \right| = \left( \sum_{k=1}^n \left| \frac{\partial f}{\partial z_k}(z) \right|^2 \right)^{1/2}.$$

Also invariant gradient of  $f$  at  $z$  defined by

$$|\tilde{\nabla} f(z)| = |\nabla(f \circ \varphi_z)(0)|.$$

The Dirichlet space  $D_\alpha^p(\mathbb{B}_n)$  is the space of the analytic functions  $f : \mathbb{B}_n \rightarrow \mathbb{C}$  such that  $R(f) \in A_\alpha^p(\mathbb{B}_n)$ . In a similar way we can define  $D_\alpha^p(\mathbb{B}_n^2)$ .

If  $f \in H(\mathbb{B}_n)$  then  $L(f) \in H(\mathbb{B}_n^2)$ . So direct calculation shows that

$$R(Lf)(z, w) = L(Rf)(z, w) + \sum_{k=1}^n \frac{(z_k - w_k)(f(w) - f(z))}{(z - w)^2}. \tag{1}$$

The operator we use here is the well-known symmetric lifting operator which lifts analytic functions from  $\mathbb{B}_n$  to  $\mathbb{B}_n^2$  and defined by

$$L : H(\mathbb{B}_n) \rightarrow H(\mathbb{B}_n^2)$$

$$L(f)(z, w) = \frac{f(z) - f(w)}{z - w},$$

where  $H(\mathbb{B}_n^2)$  is the space of all analytic functions on  $\mathbb{B}_n^2$ .

Wulan and Zhu in [8] characterized the Bergman spaces in the unit disk and unit ball in terms of Lipschitz type conditions also studied the action of operator  $L : H(\mathbb{D}) \rightarrow H(\mathbb{D}^2)$  on  $A_\alpha^p$ . Double integral characterization for  $A_\alpha^p(\mathbb{B}_n)$  can be found for example in [3, 5, 6]. In [3] the authors use the double integral characterization for proving the action of the operator  $L$  on  $A_\alpha^p$  where  $p = \alpha + 2$ . Recently, the first author of this paper and Sohrabi investigated the symmetric lifting operator on the Bloch type spaces in [2]. In this work, we investigate the action symmetric lifting operator  $L : A_\alpha^p(\mathbb{B}_n) \rightarrow A_\alpha^p(\mathbb{B}_n^2)$  and  $L : D_\alpha^p(\mathbb{B}_n) \rightarrow D_\alpha^p(\mathbb{B}_n^2)$ .

## 2. Symmetric lifting operator from $A_\alpha^p(\mathbb{B}_n)$ into $A_\alpha^p(\mathbb{B}_n^2)$

In this section we first bring some lemmas which are needed for proving the main results.

**Theorem 2.1.** [8] Suppose that  $p > 0$ ,  $\alpha > -1$ , and  $f$  is analytic in  $\mathbb{B}_n$ . Then the following conditions are equivalent.

(i)  $f \in A_\alpha^p$ .

(ii) There exists a continuous function  $g$  in  $L^p(\mathbb{B}_n, dv_\alpha)$  such that

$$|f(z) - f(w)| \leq \rho(z, w)(g(z) + g(w)), \quad z, w \in \mathbb{B}_n.$$

(iii) There exists a continuous function  $g$  in  $L^p(\mathbb{B}_n, dv_\alpha)$  such that

$$|f(z) - f(w)| \leq \beta(z, w)(g(z) + g(w)), \quad z, w \in \mathbb{B}_n.$$

(iv) There exists a continuous function  $g$  in  $L^p(\mathbb{B}_n, dv_{p+\alpha})$  such that

$$|f(z) - f(w)| \leq |z - w|(g(z) + g(w)), \quad z, w \in \mathbb{B}_n.$$

**Lemma 2.2.** ([5]) Let  $r > 0$ . Then

$$1 - |z|^2 \sim 1 - |w|^2 \sim |1 - \langle z, w \rangle|^2$$

for all  $z \in \mathbb{B}_n$  and  $w \in D(z, r)$ . Furthermore, there exists a positive constant  $C$  such that

$$(1 - |z|^2)^p |\nabla f(z)|^p \leq \frac{C}{(1 - |z|^2)^{n+1}} \int_{D(z, r)} |f(w) - f(z)|^p dv(w)$$

for all  $z \in \mathbb{B}_n$  and  $f \in H(\mathbb{B}_n)$ .

**Lemma 2.3.** [5] The involutive automorphism  $\varphi_z$  has the following properties:

$$1 - |\varphi_z(w)|^2 = \frac{(1 - |z|^2)(1 - |w|^2)}{|1 - \langle z, w \rangle|^2}$$

and

$$|\varphi_z(w)|^2 = \frac{|z - w|^2 + \langle z, w \rangle|^2 - |z|^2|w|^2}{|1 - \langle z, w \rangle|^2}.$$

Consequently

$$|\varphi_z(w)| \leq \frac{|z - w|}{|1 - \langle z, w \rangle|}$$

and

$$\frac{(1 - |z|^2)(1 - |w|^2)}{|1 - \langle z, w \rangle|^2} \leq \frac{1 - |\varphi_z(w)|^2}{|\varphi_z(w)|^2}.$$

In the next three theorems, the symmetric lifting operator  $L : A_\alpha^p(\mathbb{B}_n) \rightarrow A_\alpha^p(\mathbb{B}_n^2)$  will be considered.

**Theorem 2.4.** Suppose  $\alpha > -1$  and  $0 < p < \alpha + n + 1$ . Then the symmetric lifting operator  $L$  maps  $A_\alpha^p(\mathbb{B}_n)$  boundedly into  $A_\alpha^p(\mathbb{B}_n^2)$ .

**Proof.** Suppose that  $f \in A_\alpha^p(\mathbb{B}_n)$ . Theorem 2.1 implies that there exists a continuous function  $g \in L^p(\mathbb{B}_n, dv_\alpha)$  such that

$$|f(z) - f(w)| \leq \rho(z, w)(g(z) + g(w)), \quad z, w \in \mathbb{B}_n.$$

Since  $\rho(z, w) = |\varphi_z(w)| \leq \frac{|z-w|}{|1-\langle z, w \rangle|}$ , we have

$$\frac{|f(z) - f(w)|}{|z - w|} \leq \frac{g(z)}{|1 - \langle z, w \rangle|} + \frac{g(w)}{|1 - \langle z, w \rangle|}, \quad z, w \in \mathbb{B}_n.$$

So

$$|L(f)(z, w)|^p \leq C \left( \frac{g(z)^p}{|1 - \langle z, w \rangle|^p} + \frac{g(w)^p}{|1 - \langle z, w \rangle|^p} \right).$$

Therefore

$$\int_{\mathbb{B}_n} \int_{\mathbb{B}_n} |L(f)(z, w)|^p dv_\alpha(z) dv_\alpha(w) \leq 2C \int_{\mathbb{B}_n} g(z)^p dv_\alpha(z) \int_{\mathbb{B}_n} \frac{dv_\alpha(w)}{|1 - \langle z, w \rangle|^p}.$$

If  $p < n + 1 + \alpha$  then from Theorem 1.12 of [9] we have the boundedness of the internal integral. So

$$\int_{\mathbb{B}_n} \int_{\mathbb{B}_n} |L(f)(z, w)|^p dv_\alpha(z) dv_\alpha(w) \leq C \int_{\mathbb{B}_n} g(z)^p dv_\alpha(z).$$

We get  $L(f) \in A_\alpha^p(\mathbb{B}_n^2)$  and  $L$  maps  $A_\alpha^p(\mathbb{B}_n)$  into  $A_\alpha^p(\mathbb{B}_n^2)$ . The proof of boundedness of  $L : A_\alpha^p(\mathbb{B}_n) \rightarrow A_\alpha^p(\mathbb{B}_n^2)$  comes from the closed graph theorem. Suppose that  $(f, g) \in \overline{G}$  where  $G$  is the graph of  $L$ . Then there exists a sequence  $(f_n, Lf_n)$  of  $G$  such that  $(f_n, Lf_n) \rightarrow (f, g)$  which results  $f_n \rightarrow f$  and  $Lf_n \rightarrow g$ . One can check that using the definition of norm of the Bergman space,  $Lf_n \rightarrow Lf$ . So  $Lf = g$  which means that the graph of  $L$  is closed and the operator  $L$  is bounded.  $\square$

**Theorem 2.5.** *Suppose  $\alpha > -1$  and  $p > \alpha + n + 1$ . Then the symmetric lifting operator  $L$  maps  $A_\alpha^p(\mathbb{B}_n)$  boundedly into  $A_\beta^p(\mathbb{B}_n^2)$  where  $\beta = (p + \alpha - n - 1)/2$ .*

The proof of the above theorem is similar to the previous one by using Theorem 2.1 and Theorem 1.12[9].

**Theorem 2.6.** *Suppose  $\alpha > -1$  and  $p = \alpha + n + 1$ . Then the symmetric lifting operator  $L$  maps  $A_\alpha^p(\mathbb{B}_n)$  boundedly into  $A_\gamma^p(\mathbb{B}_n^2)$  for any  $\gamma > \alpha$ .*

**Proof.** If  $f \in A_\alpha^p$ , then by Theorem 2.1 there exists a continuous function  $g \in L^p(\mathbb{B}_n, dv_\alpha)$  such that

$$|f(z) - f(w)| \leq \rho(z, w)(g(z) + g(w)) \leq \frac{|z - w|}{|1 - \langle z, w \rangle|} (g(z) + g(w)).$$

There exists a positive constant  $C$  such that

$$\begin{aligned} \int_{\mathbb{B}_n} \int_{\mathbb{B}_n} |L(f)(z, w)|^p dv_\gamma(z) dv_\gamma(w) &\leq 2C \int_{\mathbb{B}_n} g(z)^p dv_\gamma(z) \int_{\mathbb{B}_n} \frac{dv_\gamma(w)}{|1 - \langle z, w \rangle|^p} \\ &= 2C \int_{\mathbb{B}_n} g(z)^p dv_\gamma(z) \int_{\mathbb{B}_n} \frac{dv_\gamma(w)}{|1 - \langle z, w \rangle|^{n+1+\alpha}}. \end{aligned}$$

Since  $\gamma > \alpha$ , the last integral is bounded. Then

$$\int_{\mathbb{B}_n} \int_{\mathbb{B}_n} |L(f)(z, w)|^p dv_\gamma(z) dv_\gamma(w) \leq 2C \int_{\mathbb{B}_n} g(z)^p dv_\gamma(z) < 2C \int_{\mathbb{B}_n} g(z)^p dv_\alpha(z) < \infty.$$

$\square$

### 3. Symmetric lifting operator from $D_\alpha^p(\mathbb{B}_n)$ into $D_\alpha^p(\mathbb{B}_n^2)$

In this section we first state a characterization for Dirichlet space using pseudo-hyperbolic metric and then study the symmetric lifting operator  $L$  from  $D_\alpha^p(\mathbb{B}_n)$  into  $D_\alpha^p(\mathbb{B}_n^2)$ .

The following characterization for Bergman space is crucial for our main results, [9].

**Lemma 3.1.** *Suppose that  $p > 0$ ,  $\alpha > -1$ , and  $f \in H(\mathbb{B}_n)$ . Then the following conditions are equivalent.*

- (a)  $f \in A_\alpha^p(\mathbb{B}_n)$
- (b)  $|\tilde{\nabla}f(z)|$  is in  $L^p(\mathbb{B}_n, dv_\alpha)$ .
- (c)  $(1 - |z|^2)|\nabla f(z)|$  is in  $L^p(\mathbb{B}_n, dv_\alpha)$ .
- (d)  $(1 - |z|^2)|Rf(z)|$  is in  $L^p(\mathbb{B}_n, dv_\alpha)$ .

**Theorem 3.2.** *Suppose that  $p > 0$  and  $\alpha > -1$ . If  $f \in D_\alpha^p(\mathbb{B}_n)$  then there exists a continuous function  $g \in L^p(\mathbb{B}_n, dv_\alpha)$  such that*

$$|f(z) - f(w)| \leq \rho(z, w)(g(z) + g(w)).$$

**Proof.** It can be seen that

$$f(z) - f(0) = \int_0^1 \frac{Rf(tz)}{t} dt$$

for all  $z \in \mathbb{B}_n$ . Fix  $r \in (0, 1)$ . Then we have

$$|f(z) - f(0)| \leq |z| \sup\{|\nabla f(w)| : w \in D(0, r)\}$$

where  $D(0, r)$  is the pseudo-hyperbolic disk centered at 0 with radius  $r$ . Since the Euclidean metric is comparable to the pseudo-hyperbolic metric in the relatively compact set  $D(0, r)$  and  $|\nabla f(w)|$  is comparable to  $|\tilde{\nabla} f(w)|$  in this set, there exists a positive constant  $C$  such that

$$|f(z) - f(0)| \leq C\rho(z, 0) \sup\{|\tilde{\nabla} f(w)| : w \in D(0, r)\}.$$

Put  $f \circ \varphi_w$  in place of  $f$ ,  $\rho(z, w) < r$ , and  $\varphi_w(z)$  in place of  $z$ . Then the Mobius invariancy of pseudo-hyperbolic metric and invariant gradient (see [9]) implies that

$$|f(z) - f(w)| \leq C\rho(z, w) \sup\{|\tilde{\nabla} f(u)| : u \in D(z, r)\}.$$

Set

$$h(z) = \sup\{|\tilde{\nabla} f(u)| : u \in D(z, r)\}.$$

So

$$|f(z) - f(w)| \leq C\rho(z, w)(h(z) + h(w)),$$

for  $z$  and  $w$  with  $\rho(z, w) < r$ . If  $\rho(z, w) \geq r$ , then we set

$$g(z) = \frac{|Rf(z)|}{r} + Ch(z).$$

Clearly

$$|f(z) - f(w)| \leq C\rho(z, w)(g(z) + g(w)),$$

for all  $z, w$ . It just remains that showing  $g \in L^p(\mathbb{B}_n, dv_\alpha)$ . But  $Rf \in L^p(\mathbb{B}_n, dv_\alpha)$ . So we need to show that  $h \in L^p(\mathbb{B}_n, dv_\alpha)$  which is a similar argument as in the proof of Theorem 5.1 of [8].  $\square$

**Lemma 3.3.** *Suppose  $\alpha > -1$  and  $0 < p < \alpha + n + 1$ . Then the symmetric lifting operator  $L$  maps  $D_\alpha^p(\mathbb{B}_n)$  boundedly into  $A_\alpha^p(\mathbb{B}_n^2)$ .*

The proof is the same as proof of Theorem 2.4 using the previous theorem.

In the cases  $p > \alpha + n + 1$  and  $p = \alpha + n + 1$  we have the following lemma which are stated without proof.

**Lemma 3.4.** *Suppose  $\alpha > -1$  and  $p > \alpha + n + 1$ . Then the symmetric lifting operator  $L$  maps  $D_\alpha^p(\mathbb{B}_n)$  boundedly into  $A_\beta^p(\mathbb{B}_n^2)$  where  $\beta = (p + \alpha - n - 1)/2$ .*

**Lemma 3.5.** *Suppose  $\alpha > -1$  and  $p = \alpha + n + 1$ . Then the symmetric lifting operator  $L$  maps  $D_\alpha^p(\mathbb{B}_n)$  boundedly into  $A_\gamma^p(\mathbb{B}_n^2)$  for any  $\gamma > \alpha$ .*

Now we are ready to state the main result of this section.

**Theorem 3.6.** *Suppose  $\alpha > -1$ . Then*

- (a) *For  $0 < p < \alpha + n + 1$ , the symmetric lifting operator  $L$  maps  $D_\alpha^p(\mathbb{B}_n)$  boundedly into  $D_\alpha^p(\mathbb{B}_n^2)$ .*
- (b) *For  $p > \alpha + n + 1$ , the symmetric lifting operator  $L$  maps  $D_\alpha^p(\mathbb{B}_n)$  boundedly into  $D_\beta^p(\mathbb{B}_n^2)$  where  $\beta = (p + \alpha - n - 1)/2$ .*
- (c) *For  $p = \alpha + n + 1$ , the symmetric lifting operator  $L$  maps  $D_\alpha^p(\mathbb{B}_n)$  boundedly into  $D_\gamma^p(\mathbb{B}_n^2)$  for any  $\gamma > \alpha$ .*

**Proof.** Suppose that  $f \in D_\alpha^p(\mathbb{B}_n)$ . We need to prove that  $Lf \in D_\alpha^p(\mathbb{B}_n^2)$  or equivalently  $R(Lf) \in A_\alpha^p(\mathbb{B}_n^2)$ .  $f \in D_\alpha^p(\mathbb{B}_n)$  implies that  $Rf \in A_\alpha^p(\mathbb{B}_n)$  and using Theorem 2.4, we get  $L(Rf) \in A_\alpha^p(\mathbb{B}_n^2)$ . From (1), it will be sufficient to prove that  $J \in A_\alpha^p(\mathbb{B}_n^2)$  where

$$J = \sum_{k=1}^n \frac{(z_k - w_k)(f(w) - f(z))}{(z - w)^2}.$$

Using triangle inequality and direct calculation we obtain for some positive constant  $C$

$$\begin{aligned} \int_{\mathbb{B}_n} \int_{\mathbb{B}_n} |J|^p dv_\alpha(z) dv_\alpha(w) &= \int_{\mathbb{B}_n} \int_{\mathbb{B}_n} \left| \sum_{k=1}^n \frac{(z_k - w_k)(f(w) - f(z))}{(z - w)^2} \right|^p dv_\alpha(z) dv_\alpha(w) \\ &\leq C \int_{\mathbb{B}_n} \int_{\mathbb{B}_n} \sum_{k=1}^n \frac{|z_k - w_k|^p |f(z) - f(w)|^p}{|z - w|^{2p}} dv_\alpha(z) dv_\alpha(w) \\ &= C \int_{\mathbb{B}_n} \int_{\mathbb{B}_n} \sum_{k=1}^n \frac{|z_k - w_k|^p}{|z - w|^p} \frac{|f(z) - f(w)|^p}{|z - w|^p} dv_\alpha(z) dv_\alpha(w) \\ &\leq Cn^p \int_{\mathbb{B}_n} \int_{\mathbb{B}_n} \frac{|f(z) - f(w)|^p}{|z - w|^p} dv_\alpha(z) dv_\alpha(w) \\ &= Cn^p \int_{\mathbb{B}_n} \int_{\mathbb{B}_n} |Lf(z, w)|^p dv_\alpha(z) dv_\alpha(w) \\ &< \infty. \end{aligned}$$

The last line of the above equation comes from Lemma 3.3. The proof of part (a) is completed. In the other parts we have the similar argument.  $\square$

One can see that for  $\alpha > -1$  and  $p > 0$ , the symmetric lifting operator  $L$  maps  $A_\alpha^p(\mathbb{B}_n)$  boundedly into  $A_{\alpha+p}^p(\mathbb{B}_n^2)$  and also maps  $D_\alpha^p(\mathbb{B}_n)$  boundedly into  $D_{\alpha+p}^p(\mathbb{B}_n^2)$ .

### References

- [1] P. DUREN AND A. SCHUSTER, *Bergman spaces*, vol. 100 of Mathematical Surveys and Monographs, American Mathematical Society, Providence, RI, 2004.
- [2] M. HASSANLOU AND M. SOHRABI, *Characterization of Bloch type spaces and symmetric lifting operator*, International Journal of Nonlinear Analysis and Applications, (2022).
- [3] M. HASSANLOU AND H. VAEZI, *Double integral characterization for Bergman spaces*, Iran. J. Math. Sci. Inform., 11 (2016), pp. 27–34, 148.
- [4] H. HEDENMALM, B. KORENBLUM, AND K. ZHU, *Theory of Bergman spaces*, vol. 199 of Graduate Texts in Mathematics, Springer-Verlag, New York, 2000.
- [5] S. LI, H. WULAN, R. ZHAO, AND K. ZHU, *A characterisation of Bergman spaces on the unit ball of  $\mathbb{C}^n$* , Glasg. Math. J., 51 (2009), pp. 315–330.
- [6] S. LI, H. WULAN, AND K. ZHU, *A characterization of Bergman spaces on the unit ball of  $\mathbb{C}^n$ . II*, Canad. Math. Bull., 55 (2012), pp. 146–152.
- [7] M. STESSIN AND K. ZHU, *Composition operators on embedded disks*, J. Operator Theory, 56 (2006), pp. 423–449.
- [8] H. WULAN AND K. ZHU, *Lipschitz type characterizations for Bergman spaces*, Canad. Math. Bull., 52 (2009), pp. 613–626.
- [9] K. ZHU, *Spaces of holomorphic functions in the unit ball*, vol. 226 of Graduate Texts in Mathematics, Springer-Verlag, New York, 2005.

Please cite this article using:

Mostafa Hassanlou, Ebrahim Abbasi, Bergman and Dirichlet spaces in the unit ball and symmetric lifting operator, AUT J. Math. Comput., 4(2) (2023) 155-160  
<https://doi.org/10.22060/ajmc.2022.21778.1107>

